

SURVEILLANCE OF INFECTIOUS DISEASES IN ANIMALS AND HUMANS IN SWEDEN 2019

Chapter excerpt -
Nephropathia epidemica



Editor: Karl Ståhl

Department of Disease Control and Epidemiology, National Veterinary Institute (SVA), SE-751 89 Uppsala, Sweden.

Authors: Charlotte Axén, Mia Brytting, Ioana Bujila, Erika Chenais, Rikard Dryselius, Helena Eriksson, Pernille Etterlin, Eva Forsgren, Malin Grant, Gittan Gröndahl, Gunilla Hallgren, Anette Hansen, Gete Hestvik, Marika Hjertqvist, Mia Holmberg, Cecilia Hultén, Helena Höök, Cecilia Jernberg, Jerker Jonsson, Ulrika König, Elina Lahti, Emelie Larsdotter, Mats Lindblad, Anna Lundén, Emma Löf, Hans-Olof Nilsson, Oskar Nilsson, Maria Nöremark, Anna Ohlson, Ylva Persson, Karin Persson-Waller, Thomas Rosendal, Karl Ståhl, Robert Söderlund, Kaisa Sörén, Magnus Thelander, Karin Troell, Anders Wallensten, Per Wallgren, Stefan Widgren, Ulrika Windahl, Beth Young, Nabil Yousef, Siamak Zohari, Erik Ågren, Estelle Ågren, Elina Åsbjer

Cover Photo: Anders Lindström

Copyright of map data: ©EuroGeographics for the administrative boundaries

Reporting guidelines: Reporting guidelines were introduced in 2018 for those those chapters related to purely animal pathogens. The guidelines build on experiences from several EU projects, and have been validated by a team of international experts in animal health surveillance. The aim is to develop these guidelines further in collaboration within the global surveillance community and they have therefore been made available in the form of a wiki on the collaborative platform GitHub (<https://github.com/SVA-SE/AHSURED/wiki>). Feel free to contribute!

Layout: The production of this report continues to be accomplished using a primarily open-source toolset. The method allows the source text, produced by authors, to be edited independently of the template for the layout which can be modified and reused for future reports. Specifically, the chapter texts, tables and captions are authored in Microsoft Word and then converted using pandoc and R to the LaTeX typesetting language. Most figures and maps are produced using the R software for statistical computing. Development for 2019 has further improved the importing of content from Word to LaTeX. The method can now import text, tables and figure captions from Word, as well as the newly designed 'IN FOCUS' sections of some chapters. The tool is available as an R-package at GitHub (<https://github.com/SVA-SE/mill/>). This year the report was also built with a continuous integration pipeline on Microsoft's Azure DevOps platform, allowing every committed change to the content to be built and tested automatically. The report generation R-package and process was designed by Thomas Rosendal and Stefan Widgren. In 2019, figures and the final typesetting were done by Wiktor Gustafsson and Thomas Rosendal with contributions from the report authors.

Print: TMG Tabergs AB.

Text, tables, figures and maps may be cited and reprinted only with reference to this report.

Suggestion citation: Surveillance of infectious diseases in animals and humans in Sweden 2019, National Veterinary Institute (SVA), Uppsala, Sweden. SVA:s rapportserie 64 1654-7098.

This report may be subject to updates and corrections. The latest version is always available for download at www.sva.se.

Nephropathia epidemica



Humans may be exposed to Puumala virus during occupational or recreational activities, such as working with hay, cleaning barns or summer cottages, cutting wood and entering buildings contaminated with rodent excretions. Photo: Butivshchenko Olena/Shutterstock.

BACKGROUND

Nephropathia epidemica (NE) is caused by Puumala virus, a member of the Hantavirus genus in the *Bunyaviridae* family. Hantaviruses are the cause of rodent-borne haemorrhagic fevers with renal syndrome (HFRS) and hantavirus pulmonary syndrome (HPS). Puumala virus is likely to be the most prevalent hantavirus in Europe. The virus is excreted in saliva, urine and faeces from its natural reservoir, the bank vole. Puumala virus can remain infectious in bank vole cage bedding for two weeks. Transmission to humans often occurs in an aerosolised form. Humans may be exposed to virus aerosols during occupational or recreational activities, such as working with hay, cleaning barns or summer cottages, cutting wood and entering buildings contaminated with rodent excretions.

Nephropathia epidemica was first described by two Swedish physicians, independently, in 1934. The linkage to the bank vole was suggested many years later. The virus was first isolated in 1982 in Puumala, a municipality in south-eastern Finland.

In Sweden, between 50 and 600 cases are reported each season with a considerable interannual variation coupled to the 3–4-year population cycle of the bank vole. During the winter seasons 2006–2007 and 2007–2008 the number of

notified cases rose to 1400, where most of the cases occurred in 2007 (Figure 14). It is hypothesised that a parallel occurrence of a peak in the bank vole population and lack of snow cover in December 2006 caused bank voles to seek refuge in buildings and barns, hence increasing their contact with humans.

DISEASE

Animals

In the bank vole, the infection is understood to be subclinical.

Humans

The clinical picture is characterised by a sudden onset of high fever, headache, backache and abdominal pain. The symptoms range from sub-clinical to renal failure requiring intensive care and dialysis, but fatal cases are rare. The incubation period varies from 2 to 6 weeks.

LEGISLATION

Animals

Hantaviruses are not notifiable in animals.

Humans

Nephropathia epidemica has been notifiable since 1989 according to the Communicable Disease Act (SFS 2004:168 with the amendments of SFS 2013:634).

SURVEILLANCE

Animals

There is no surveillance in animals.

Humans

The surveillance in humans is mandatory and based on identification of the disease by a treating physician or by laboratory diagnosis. Both are obligated to report identified cases to the regional and national level to enable further analyses and adequate intervention measures.

RESULTS

Humans

In 2019, 155 cases of NE were reported, which was a decrease in comparison to the previous year (Figure 14). The median age among all cases was 57 and most reported cases were males in the age category 25 years and older. There were very few cases below the age of 25 years reported, both among men and women. Consistent with previous years, more cases were reported in men (63%) than in women. The reason for this difference in incidence between age groups and sexes is not completely understood, but behaviour is

most likely an important factor.

Most of the reported NE cases acquire their infections in Sweden. In 2019, there was only one case who had been infected in Finland as well as fifteen cases with unknown country of infection.

A majority of the cases were reported to have been infected in Norrland and the northern parts of Svealand. The incidence was highest in the County of Norrbotten (24 cases per 100 000 inhabitants) followed by the County of Västerbotten (18 cases per 100 000 inhabitants). All cases reported from the southern parts of Sweden were infected further north, i.e. in areas where NE is already known to occur. This regional pattern is consistent with patterns observed during previous years.

DISCUSSION

During recent years, fluctuations in the bank vole population have coincided with increases and decreases in the number of human cases of Puumala virus infections. The 3–4-year natural population cycle and variations in the climatic conditions impact the rodent populations.

REFERENCES

Evander M, Ahlm C (2009) Milder winters in northern Scandinavia may contribute to larger outbreaks of haemorrhagic fever virus. *Global Health Action* 2:10.3402/gha.v2i0.2020




Figure 14: Notified incidence per 100 000 inhabitants of human Nephropathia epidemica in Sweden 1998–2019.